

# 2

## **Cervical Cancer Screening Strategies for Aboriginal Women<sup>1</sup>**

Agnes T. Black<sup>2</sup>

### **Introduction**

British Columbia is home to the British Columbia Cancer Agency's Cervical Cancer Screening Program (CCSP), a population-based cervical cancer screening program credited with achieving a 70% reduction in cervical cancer incidence in British Columbia over the past fifty years, largely by widely promoting screening and making screening services readily available and free of cost (BC Cancer Agency 2007). While the CCSP has had enormous success in increasing rates of cervical cancer screening among most of the province's female population, Aboriginal women in British Columbia are estimated to have Pap rates of 50% compared to 85% for all women in British Columbia (Hislop et al. 1996). Mortality from cervical cancer is six times higher among Aboriginal women than other women in British Columbia (Hislop et al. 1996).

The CCSP does not collect data on the racial or ethnic background of women participating in cervical cancer screening, and current rates of Pap screening among Aboriginal women are difficult to determine from other available data sources. However, practitioners strongly suspect that Aboriginal women are not participating in cervical cancer screening in the same numbers as their non-Aboriginal counterparts. This project was started based on the hypothesis that Aboriginal women are not participating in cervical cancer screening in any greater percentages than they were when Hislop did an analysis in 1996.

Previous qualitative research studies have examined reasons for the reduced rate of screening among Aboriginal women, and recommended interventions aimed at improving screening rates among this population (Clarke et al. 1998; Pakula 2006). This research project built on previous studies and explored the question: "What interventions currently exist, or have been proposed, to encourage participation in cervical cancer screening among Aboriginal women, and what are the barriers and facilitators to implementing these interventions?" We conducted an interview-based study focused on awareness and evaluation of current interventions to encourage cervical cancer screening among Aboriginal women.

The objective of this research was twofold: To provide the CCSP with an evaluation of the current state of cervical cancer screening interventions for Aboriginal women in British Columbia, and to offer suggestions for networking and building relationships with Aboriginal communities in British Columbia and organizing

outreach efforts in these communities. This information will assist the CCSP in choosing and tailoring future interventions to improve cervical cancer screening rates for Aboriginal women.

This objective was accomplished by the completion of sixteen key informant interviews, along with eleven surveys, the results of which are found in this report. It is hoped that the recommendations found in this report will further the goals of the CCSP, which include reducing morbidity from cervical dysplasia (the precursor to cervical cancer), and reducing mortality from cervical cancer. The CCSP has initiated primarily quantitative research projects, therefore this project presented an opportunity to demonstrate the effectiveness and usefulness of qualitative research findings in supporting the work and the goals of the CCSP.

## Methodology

**Study Design.** This project involved exploratory research, approached through qualitative research methods that emphasize developing insights and generalizations from the data collected (Neuman 2006). Site visits in different parts of the province allowed the researcher to gather other information for use in triangulation,<sup>3</sup> and therefore benefit from studying cervical cancer screening interventions from multiple points of view. The primary methodological tool employed in this project was key informant interviews, and a total of sixteen people were interviewed. A survey was also used at one site visit.<sup>4</sup>

**Interviewees.** The people interviewed for this project included physicians, nurses, health authority staff, band office staff, and others working in the field of Aboriginal women's health. Interviews were sought with people working with groups of Aboriginal women, rather than interviews with individual Aboriginal women. This decision was made in part because previous research had focused on individual barriers to participation in cervical cancer screening, while this project focused on organization barriers and facilitators to participation. It was also easier to identify and procure access to people at the clinic, band, or health authority level, rather than locating individual Aboriginal women who would agree to be interviewed by a health-care researcher.

Eight of the sixteen people interviewed self-identified as Aboriginal, and their responses to interview questions sometimes included personal reflections on being an Aboriginal woman in the health-care system.

A survey was designed and used at one point in the project, when the researcher was invited to attend a day-long meeting of the Fort St. John Aboriginal Health Improvement Committee (AHIC). Because time constraints did not permit individual interviews with each of the committee members, a brief survey was distributed with the permission of the meeting's facilitator, and members of the committee were encouraged to fill out the survey. The results of this survey are included in the results section of this report.

**Study Site and Population.** A purposive sample of several geographical areas was chosen to include both urban and rural regions. Previous research has shown that barriers and facilitators to cervical cancer screening differ among urban and rural Aboriginal women. Areas included in the project were Vancouver, Victoria, Port Alberni, Nanaimo, and Fort St. John.

**Sampling Plan.** Key informants in each of the geographical areas were identified through snowball sampling. Key informants consisted of a purposive sample of sixteen people, including health authority staff working in the area of Aboriginal women's health, health coordinators hired by First Nations bands, and health professionals providing cervical cancer screening to Aboriginal women. Surveys were completed by eleven members of the Aboriginal Health Improvement Committee in Fort St. John.

**Data Collection Instruments.** Interviews consisted of open-ended questions to encourage participants to offer opinions and propose solutions. Most interviews were tape-recorded, although two interviews were transcribed from handwritten notes, at the preference of the interviewees. At three different times, two women were interviewed at a time, at their preference.

**Procedures.** Informed consent was received from each participant prior to beginning interviews. The names of interviewees were changed to pseudonyms, and tapes were then transcribed and analyzed.

In Victoria and Vancouver, key informants were identified through snowball sampling, asking for potential interviewee names from others working in the field of Aboriginal women's health. Attempts were made to include community health representatives (CHRs) for First Nations bands, but no CHRs were available to participate, and some expressed the opinion that they had previously participated in research projects but had not seen benefits to their patients or their bands.

Site visits were conducted in Victoria, Port Alberni, Vancouver, and Fort St. John. When in-person interviews could not be scheduled, telephone interviews were conducted with interviewees in Nanaimo, Port Alberni, and Victoria.

**Analysis Plan.** Transcriptions of interviews were thematically coded to identify specific barriers and facilitators. Open coding was performed, followed by axial coding to further organize ideas and themes emerging from the interview transcriptions (Neuman 2006). Data was coded using Microsoft Word, and results are described in this report.

**Limitations of the Study.** This study is restricted to several geographical areas in British Columbia, which limits the degree to which the results and recommendations can be generalized.

The perspectives of individual Aboriginal women accessing the health-care system are not included, as interviews were conducted solely with practitioners and administrators in the health-care field who worked with Aboriginal women; therefore, this study does not encompass the full range of opinions held by all Aboriginal women in British Columbia.

**Ethical Considerations.** Ethics approval was obtained from the Simon Fraser University Research Ethics Board and the University of British Columbia—British Columbia Cancer Agency Research Ethics Board prior to beginning interviews for this study. All protocols were observed.

As a non-Aboriginal person, I recognized the inherent challenges in developing relationships and trust in the Aboriginal communities I visited and with the Aboriginal women I interviewed, especially considering the short time frame of this project. I relied on the mentorship of others who have worked with Aboriginal women or done research in this field, and followed their advice on partnership building and respectful approaches. I encouraged the people I interviewed to describe their population's health challenges and priorities, so that my priorities as a researcher would not dominate the interview. I remain committed to full distribution of the results of this research in order to benefit the Aboriginal women whom the project was designed to serve.

**Knowledge Exchange.** Copies of this report were distributed to each of the sixteen interviewees, and to the facilitator of the AHIC meeting at which the surveys were distributed. Additional copies will be made and distributed by the CCSP as needed or desired. The results will be presented to the CCSP.

## **Current Interventions to Increase Aboriginal Women's Participation in Cervical Cancer Screening**

There are several programs in British Columbia that are dedicated to improving the health of the Aboriginal women in the province. Most of these programs give some attention to cervical cancer screening.

All health authorities in British Columbia have Aboriginal health plans, programs, or teams, with staff dedicated to health improvement in Aboriginal communities. Within the Provincial Health Services Authority (PHSA) there is an Aboriginal Health Program at BC Women's Hospital and Health Centre that provides outreach, health education, and cervical cancer screening in remote reserves. This program serves between 150 and 200 Aboriginal women per year.

Also included in PHSA is the BC Centre for Disease Control, which contains the Street Nurse Program (SNP). The nurses of the SNP have been organizing "Papalooza" in Vancouver's Downtown Eastside since 2002. The nurses perform between fifty and one hundred Pap tests during this three-day event, which is hosted once or twice a year. Up to 42% of the participants in Papalooza have been Aboriginal women (Taylor, personal communication, June 11, 2008).

The Vancouver Island Health Authority (VIHA) developed the Aboriginal Women's Wellness Program. This program has a coordinator whose work includes women's health promotion and outreach, and a family nurse-practitioner whose work also includes women's health promotion and outreach, as well as cervical cancer screening for the local Aboriginal population.

Vancouver Coastal Health has an Aboriginal Health Initiative Program that supports health promotion projects with Aboriginal and First Nations organizations within the Vancouver Coastal Health region. Fraser Health Authority has an Aboriginal Health Initiatives Program and has just hired a nurse-practitioner to serve Aboriginal clients at the Kla-how-eya Aboriginal Centre Health Clinic in Surrey. Interior Health Authority and Northern Health Authority each have an Aboriginal health program or team.

Within the health regions there are individual efforts to increase cervical cancer screening rates among Aboriginal women. An example of this occurred recently in Port Hardy, where the public health nurses were funded by the health authority to organize, advertise and host a “Pap Week” event, which included nurse-run clinics performing Pap tests in eight Aboriginal communities. In all, this week-long series of clinics served three hundred women and performed sixty-nine Pap tests (Rardon, personal communication, June 17, 2008).

There are federal health nurses in nineteen communities across British Columbia, employed by Health Canada with a mandate to provide health services to the on-reserve First Nations population in their communities. This has often included support for cervical cancer screening on local reserves.

Many individual First Nations bands and tribal associations have health directors and staff dedicated to health promotion activities in their communities. In many cases, these health promotion activities include providing cervical cancer screening.

## **Interview Results**

The results of the sixteen interviews and eleven surveys are grouped into three categories: background health challenges for Aboriginal women; barriers to participation in cervical cancer screening; and outlines of interventions that are currently working to increase Aboriginal women’s participation in cervical cancer screening, or interventions that are suggested to increase participation.

The background challenges facing many of the Aboriginal women served by the people interviewed include the effects of colonialism, which are perhaps most evident in the persistent poverty facing many Aboriginal people. These social determinants of health often promote poor health and prevent participation in health promotion programs like Pap tests. Dennis Raphael’s recent research confirms that “living conditions are the most important factors shaping health and predicting the onset of disease and premature mortality” (2008), while Byers and colleagues recently reported that cancer mortality is 35% higher among patients with a lower socio-economic status (Gandey 2008).

The CCSP cannot eliminate the problems faced by those members of the population suffering the ill health effects of poverty, but it is important that these conditions be acknowledged as barriers to participation in cervical cancer screening. Interventions designed to increase Aboriginal women’s participation in cervical

cancer screening must take into account the challenges faced by women living in poor social conditions and tailor programs accordingly.

The results of the interviews and surveys illustrate these and other health challenges and discuss how they interfere with the participation of some Aboriginal women in cervical cancer screening.

### ***Background Health Challenges***

When asked to describe the background challenges, or social determinants, of the health of the Aboriginal women served by themselves or their programs, interviewees gave a number of responses. The primary challenge forming a backdrop to all other more specific concerns is the issue of poverty, and the health challenges that often accompany poverty. When questioned about health challenges of the women in her community, one interviewee hesitated, then said, “I think a lot of these women are worried about putting healthy food on the table for their kids.”

Specific health conditions and social conditions affecting Aboriginal women’s health were named by interviewees to include mitigating factors such as poverty, isolation, stress, need for healthy relationships, high drop-out rates from secondary school, high unemployment rates, domestic violence, and food insecurity; diseases such as alcoholism, drug abuse, arthritis, and diabetes; and lack of consistent access to health-care providers.

One interviewee explained, “I think the health concerns of women, like the health concerns of all our community, stem back to some of the problems of colonization and residential schools. So it’s a lot of historical stuff that we’re dealing with still today and we see that in the health issues that we deal with.”

The responses I received in interviews and surveys make it clear that the social determinants of health, including poverty and the legacy of colonialism, are major influences on the health of Aboriginal women.

### ***Barriers to Aboriginal Women Participating in Cervical Cancer Screening***

I asked interviewees to describe the barriers that prevent Aboriginal women from participating in cervical cancer screening. Survey respondents were given a choice of potential barriers and asked to mark those barriers they felt were relevant. Interview and survey responses are organized into the following themes: poverty, lack of cultural safety, legacy of paternalism, privacy and confidentiality concerns, Pap tests not seen as a priority, history of sexual abuse, availability of health-care providers, and fear of focusing on cancer.

#### ***Poverty***

Many of the women I interviewed mentioned persistent poverty, and the social challenges that coexist with poverty, as barriers to Aboriginal women’s participation in cervical cancer screening. Chief among these social challenges are the daily struggles for quality child care, adequate employment, and reliable

transportation, all of which affect women's ability to participate in cervical cancer screening.

Lack of transportation to health-care providers and facilities to participate in Pap tests was listed as a barrier by almost every person interviewed, including one who named access as a "huge issue," especially for women living in rural areas or on remote reserves.

Two-thirds of survey respondents felt that transportation was a barrier to Aboriginal women's participation in cervical cancer screening, while 27% felt that child care was a barrier. Both barriers coexist with poverty.

One interviewee stated, "A lot of women are leading very chaotic lives for a lot of reasons, and their self-care is not on their radar." Another interviewee mentioned that "The barriers are about actually having the support to take care of yourself and getting the support from other women and their families or the communities to help you do that."

### ***Lack of Cultural Safety***

The term "cultural safety," and the issues of trust and respect that are embedded in the concept of cultural safety, were mentioned by many of the women interviewed. Cultural safety is a concept that originated in New Zealand as a response to the effects of colonization on both Aboriginal peoples and on health-care professionals' treatment of Aboriginal clients. Cultural safety has been defined in many ways. Maori nurse Irihapeti Ramsden is considered the originator of the concept, and she writes, "Cultural safety addresses quality in healthcare though issues of communication and access to the health service" (Papps and Ramsden 1996). Employed as both a theoretical lens through which to consider and evaluate interactions between the health-care system and Aboriginal peoples, and a competency towards which health-care professionals can work, cultural safety has also been described as not simply cultural sensitivity, but also "an analysis of power imbalances, institutional discrimination, and the nature of relationships between the colonized and the colonizers as they apply to health care interactions" (Browne, Fiske, and Thomas 2000, 8–9). One interviewee expressed the barrier in this way: "Just having people culturally sensitized and having cultural safety, for Aboriginal people to seek help when they suspect a problem ... can be an obstacle." Another interviewee noted that barriers include "the stereotypes, the discrimination, the racism. I think those are the major challenges we have to overcome with Aboriginal people."

A health professional spoke of the lack of trust that often interferes with Aboriginal women seeking health care: "A lot of people have experienced some really negative interactions with non-First Nations people, and that brings ... a certain level of lack of trust and caution in dealing with people, so that also makes them reluctant to just go out there and keep trying. If they have a bad encounter with one physician or one clinic, they're not going to just keep doing it ... So they just stop. What happens is that they wind up using 'emerg' or walk-in clinics as

their primary care, but they fall through the cracks because neither of those places do primary care.”

### ***“Legacy of Paternalism”***

One of the health professionals I interviewed spoke of the health effects from what she called “the legacy of paternalism.” She described the ways in which her Aboriginal clients had been encouraged by health professionals in their communities to allow others to make health-care decisions for them: “In order to encourage people to use the health-care services provided by the government through the last century, they have undermined the individual’s decision-making and ownership of their health ... So many people from a First Nations community ... may have experienced a lack of independence in their own health care.” This interviewee went on to state that many of her female patients wait to be told that they need a Pap test, and in the absence of reminders, they don’t participate in cervical cancer screening.

### ***Privacy and Confidentiality Concerns***

Several of the women I interviewed noted that “shyness” and privacy were barriers for Aboriginal women participating in Pap tests. One interviewee stated that Pap tests are considered private because they are related to sexual activity, therefore there is some hesitance to discuss or undergo the Pap test. Sixty-four percent of survey respondents felt that Aboriginal women’s discomfort with the Pap test is a barrier, although respondents were not asked whether this discomfort was related to the invasiveness of the test.

In one interview, the respondent noted that women living on-reserve have told her they don’t want to go for a Pap test when the health professional visits the reserve, because “everyone will know my business” if they see her showing up for an appointment with the health professional. In another interview, concerns about confidentiality arose when discussing the training of local nurses to perform Pap tests: “When we ask them [women in community] about whether or not they would support their own community health nurse providing the services [Pap tests], they say no ... They want that confidentiality.”

Regarding confidentiality in smaller communities, one interviewee said, “It’s like any community, it’s not just Aboriginal communities. It’s all small communities. ‘Oh I seen so-and-so go down to the doctor, what’s goin’ on?’” Another interviewee stated, “I know a lot of First Nations communities who are interested in anonymity and privacy, and they don’t feel they get it with the on-reserve nurse.”

An issue related to confidentiality was described in this way by an interviewee: “There are practitioner issues. The nurse in the community can be seen, for instance as similar to child services ... They may be involved in taking a child into custody, and then you’re supposed to go to this same nurse to do your well woman exam?”



This issue was countered by another interviewee, who stated that in her community the local nurse “developed a trusting relationship with the client, and so they were quite happy with ... [the local nurse performing their Pap].”

### ***Pap Test Not Seen as a Priority***

“Paps are not on the radar, really, as a priority,” stated one interviewee when asked to describe barriers. Other interviewees explained that many Aboriginal women have other, more pressing, concerns: “They would say ‘no, my children are my priority, and my elders are my priority, my youth are my priority.’”

Some interviewees also noted a lack of awareness of the importance of cervical cancer screening, and knowledge about what the Pap test is: “I don’t think they all know what a Pap test does anymore.”

### ***History of Abuse***

The majority of interviewees mentioned sexual or physical abuse as a barrier to Pap screening among some Aboriginal women, sometimes noting that this abuse occurred in residential schools. The option of having the Pap test performed by a female provider may overcome this barrier for some women, but as one interviewee commented, “For any woman who has experienced sexual abuse ... a pelvic examination is a no-go zone, regardless of who’s doing it.”

### ***Accessibility of Health-Care Providers***

Accessibility of health-care providers was described as a barrier in those areas where there are not enough physicians or nurses. One interviewee described this situation: “So we actually just got invited to ... do a clinic and it was with the support of the physicians because the physicians are so overworked right now they can’t do just the routine (Pap) screening. They can’t get enough ... office time.” Of the survey respondents, 36% felt it was difficult for Aboriginal women to find providers taking new patients, and 64% felt it was difficult to find a female health-care provider. Access was not always seen as an issue, and, in one smaller city, a health provider stated, “If they don’t mind seeing the new doctor in town, they can have a Pap today.”

### ***Fear of Focusing on Cancer***

One interviewee stated that some of the Aboriginal women she works with have a hesitancy to discuss cancer or cancer risks because of a belief that “what you focus on, you get.” Another interviewee noted, “I know a lot of them (Aboriginal women) are really intimidated, because they’re frightened about what they’ll find ... The word ‘cancer’ just terrifies everyone.” Forty-five percent of survey respondents felt that fear of focusing on cancer was a barrier to Aboriginal women’s participation in cervical cancer screening.

In summary, the barriers preventing many Aboriginal women from participating in cervical cancer screening are primarily those related to poverty, lack of

accessibility to health-care providers, and lack of culturally safe relationships with health professionals and the health-care system.

## **Suggestions to Improve Numbers of Aboriginal Women Participating in Cervical Cancer Screening**

When the researcher discussed suggestions for interventions to increase Aboriginal women's participation in cervical cancer screening, the answers fell into several categories:

1. Build partnerships in the communities of the women you're trying to reach
2. Educate women from a young age about the importance of cervical cancer screening and the importance of the HPV vaccine in preventing cervical cancer
3. Create brochures or other educational material that reflect Aboriginal women's lives
4. Organize a day or week dedicated to Pap screening
5. Bring services to the women who live in remote communities
6. Offer drop-in appointments
7. Use creative technology for hard-to-reach populations

Below are details on each of these suggestions, followed by notes on which barriers could be overcome by each of the suggested interventions. **Table 2.1** (page 28) shows a comparison of barriers and corresponding interventions.

### ***1. Build partnerships in Aboriginal communities with individuals, CHRs, agencies, and health-authority staff working with this population.***

Many of the health-authority staff whose work currently includes a focus on Aboriginal women's health indicated that they would like to strengthen their connection to provincial health agencies. These staff people organize health conferences, visits to reserve, and Pap events, all focused on Aboriginal women's health. Participating in some of these events would provide networking opportunities that would benefit both provincial health agencies and the event organizers. Provincial health agencies could also provide materials to enhance the training in cervical cancer screening currently received by CHRs.

Several interviewees who had previously participated in Pap outreach efforts stressed the importance of time invested in building relationships with communities. One interviewee stated, "I'll come back a year later and they'll remember me ... so maybe in a small town that relationship is more than knowing them, it's also about trust and a reputation of things that get passed on ... what I notice is that our numbers will go up over time. You get that, 'Oh, they came and they weren't so bad.'"

In Port Hardy, the nurses who organized a Pap week event this year were told by the local band nurses that more women have stated they will attend the Pap week event next time because “now they know the nurses.”

Building respectful partnerships in Aboriginal communities with individuals, CHRs, agencies, and health-authority staff working with this population would help overcome the barriers presented by poverty and lack of cultural safety. This can best be accomplished by actively listening to the community and their expression of health-care priorities, then working with leaders and women’s health champions in the community to set health improvement goals that are mutually agreeable. It is important that researchers approaching Aboriginal communities be aware of the history of colonialism and the far-reaching impacts of colonialism on Aboriginal people’s health today.

***2. Educate women from a young age about the importance of cervical cancer screening, and the importance of the HPV vaccine to prevent cervical cancer.***

Several interviewees suggested that young women should be educated about the importance of cervical cancer screening and prevention, including a focus on the HPV vaccine and the Pap test. “Normalize the Pap test,” stated one health provider. Seventy-three percent of survey respondents supported increasing cervical cancer prevention education provided to Aboriginal women to encourage participation in cervical cancer screening.

One interviewee suggested that many communities seem more willing to accept educational interventions that target young people: “It gets accepted in the community easier if it’s under the rubric of ‘our youth need this.’”

Another interviewee suggested that the promotion of the HPV vaccine should be a primary goal for cervical cancer prevention efforts:

If I were to focus my efforts in cervical cancer prevention for Aboriginal communities, I would focus it right now on the HPV vaccine. Because I still think that we have monstrous health inequities due to isolation, due to access to care ... When we look at who shoulders the burden of cervical cancer in Canada, there’s the socio-economic disparity, there’s Aboriginal women, and immigrants. And for two out of three of those groups with a vaccine program that’s publicly funded and available through school-based programs, we’re going to take a big bite out of that.

Educating women from a young age about the importance of cervical cancer screening, and the importance of the HPV vaccine to prevent cervical cancer would help to overcome the barrier of the legacy of paternalism by encouraging the empowerment of young Aboriginal women to take control of their own health.

This education would also work to overcome the barriers of non-prioritization of Pap tests and fear of focusing on cancer, by combating misinformation about cervical cancer and stressing the benefits of prevention and screening.

### ***3. Create brochures and other educational material that reflect Aboriginal women's lives.***

One interviewee stated that educational material must be “reflective of the women that you’re trying to promote this screening to ... not just the young women. It’s got to be older women as well.” Another interviewee suggested that educational materials about cervical cancer screening should focus on the benefits to women and their families of being healthy: “One thing I always begin with, with the group, is that if you’re healthy then you can ... continue to care for your children. And we know it as mothers, right? We start with the women’s health, because if she has healthy children then you’ll have a healthy community ... If every woman was healthy then she would have healthy children.”

Interviewees suggested that educational material and programs should include Aboriginal role models: “To get women doing it [participating in Pap tests], you need to have First Nations role models involved. They need campaigns. They need advertisements, and explaining why it’s important for their beliefs.” Another interviewee suggested that we “tell success stories” of Aboriginal women who have been screened for cervical cancer and treated successfully, thus preventing mortality.

One suggestion was that brochures and other educational material should include other women’s faces along with Aboriginal women’s faces: “Paps are low among three groups. It’s not just Aboriginal women. Aboriginals, young women, and immigrant women, and I kind of think it’d be nice if somehow a resource could be developed with all three in mind ... One of the reasons is I don’t like the targeting like, right away you think okay, yeah, here we are, we are targeted with this again, like we’re the ones who don’t get Pap tests.”

Education about cervical cancer screening should be placed in the context of overall health and well-being, suggested one interviewee. She said health providers should make the Pap clinic be “the place in the community ... where you do the wellness, and then encompassing it [the Pap test] in wellness, it’s not just come in and have a few cells taken from your cervix, but we care about your whole health and well-being.”

Creating brochures and other educational materials that show the faces of Aboriginal women and include text that is relevant to and appropriate for Aboriginal women, with a focus on overall well-being, would assist in overcoming the barriers of cultural safety (by demonstrating respect for culture); the barrier of paternalism (by encouraging the empowerment of Aboriginal women to take control of their own health-care needs); and non-prioritization of Pap test (by combating misinformation about cervical cancer and stressing the benefits of prevention and screening).

#### ***4. Organize a day or week dedicated to Pap screening.***

Several of the interviewees for this project had previously participated in organizing events like “Pap Week” or “Papalooza,” well-publicized events with Pap tests available on a drop-in basis, and sometimes door prizes, flowers, or other incentive gifts to encourage participation and contribute to the positive atmosphere of the event. One interviewee explained to me that Papalooza is successful because “it’s a celebratory thing. It’s available in the women’s neighbourhood, and we fuss over them.”

Several interviewees told me that the CHRs in their community or reserve can assist with advertising and recruiting women to participate in a Pap day or Pap week event. Fifty-five percent of survey respondents felt that a Pap Day event would increase Aboriginal women’s participation in cervical cancer screening.

While most interviewees were enthusiastic about dedicated days or weeks, there were two cautionary notes expressed. One is the difficulty for providers in deferring all other patient concerns in order to focus on the Pap test in a short appointment time. One provider expressed her caution this way: “I think, how can you ask a question and then when they tell you the answer go, ‘Thank you!’ and carry on? ‘Have you ever been abused?’ or those invasive things, and they say ‘yes.’ They tell you their health issues and you go, ‘That was really nice. I’ve noted that. Now we’re moving on to talk about your depression,’ or something. I actually find it hard ... to speed up.”

The second caution was about confidentiality at well-publicized Pap events. One nurse stated that CHRs have told her that some Aboriginal women won’t show up at a dedicated Pap week event because they fear that other members of their community will see them at the event and their privacy will be compromised.

Having a dedicated Pap day or Pap week would help overcome the barrier of lack of availability of health-care providers, especially in remote reserves.

#### ***5. Bring Pap tests to Aboriginal women living in remote communities.***

Several interviewees stated that the best approach to improving cervical cancer screening rates in Aboriginal communities would be to deliver Pap tests in the community, performed by female providers who have established a relationship with the community, and have gained the trust of the women in the community. Fifty-five percent of survey respondents supported the approach of bringing cervical cancer screening to Aboriginal women living in remote communities.

One interviewee cautioned that Pap day or Pap week events should become a regular feature, rather than a one-time event: “One-offs are not a good idea, because the women feel like they’re being researched, then forgotten.” Another interviewee emphasized that “consistency makes a big difference ... People start relying on it and knowing it’s there so ... it has to be consistent.”

Bringing cervical cancer screening to Aboriginal women living in remote communities would help to overcome the barriers of poverty, lack of transportation, and lack of child care.

Some interviewees expressed caution about using the nurse who is from the community they're serving. One interviewee commented, "So there may be an issue of, a little bit of distance is a good thing, that it's a trusted provider but not a provider who lives right in the community." Other interviewees said the opposite, noting that in their communities the local nurse was a trusted provider and they had heard no concerns about privacy.

One interviewee expressed her experience with the difficulty of retaining nurses who have been trained to perform Paps in the reserve where they reside: "So there's the barrier of actually having practitioners. A lot of real issues, particularly on-reserve, is this cycling through of nurses. We looked over the past ... eight to ten years, and we had trained ninety nurses to do STI testing ... Five were still practicing ... There's no ability to retain nurses. It's very tough work ... the reserves are very isolated."

Her concern was echoed by another interviewee who expressed concern about the cost to the individual band of training nurses to perform Paps: "There was a lot of time and effort put into ... training them ... Those that were trained soon left the community. So the communities didn't benefit from their ... professionals taking this training. So it was costly for them, they'd have to pay for their nurses to leave the community, accommodations, travel, everything, their wage."

Another concern was expressed regarding these nurses having the support of local physicians to whom they can refer patients when the Pap test is abnormal, or when there are other issues beyond the nurses' scope of practice: "One of the things we asked each nurse before they came down [for the Pap training] is whether or not they had the support of the community, and the support of a local physician, because when you see something and it needs to be referred on to a physician you need to have that good working relationship with the physician. We found over the years that some physicians, said no, 'We offer this service. Why should we support the nurses to offer this in their community? People can come in to see us.'"

By bringing in health-care professionals from outside the reserve community, the concern about confidentiality can be addressed, as well as the concerns about retention of on-reserve nurses and costs to individual bands to train nurses. Health professionals coming to reserves to offer cervical cancer screening should work to establish collegial relationships with local health-care providers and others working locally on Aboriginal women's health issues, so they are seen as helpful and not adversarial.

### **6. Offer drop-in appointments.**

Several of the health professionals I interviewed noted that having drop-in appointments available seemed to work well for their Aboriginal clients,

especially those clients for whom lack of transportation and lack of child care were barriers to participation in cervical cancer screening. Forty-five percent of survey respondents felt that having drop-in appointments available for Pap tests would encourage participation of Aboriginal women in cervical cancer screening. Offering drop-in appointments would help address the barriers of lack of transportation and child care, as women could access health services on their own schedule, at times when transportation and child care are available to them.

***7. Use creative technology for hard-to-reach populations.***

One interviewee noted a recent research study done in Vancouver that enlisted women at high risk for HPV infection (and thus cervical cancer) to participate in cervical cancer screening by performing “self-collection” of cervical cells. (Ogilvie et al. 2007). The study addressed the need for creative approaches for women who do not participate in cervical cancer screening at recommended intervals. This intervention would help with overcoming the barrier of sexual abuse, as there is no pelvic examination to undergo, and the woman herself is in control of the specimen collection.

**Table 2.1: Barriers and Interventions**

<b>Barrier</b>	<b>Intervention</b>
Poverty, including reduced access to transportation and child care	<ul style="list-style-type: none"> <li>● Build partnerships in communities of Aboriginal women</li> <li>● Bring services to the women who live in remote communities</li> <li>● Offer drop-in appointments</li> </ul>
Lack of cultural safety	<ul style="list-style-type: none"> <li>● Build partnerships in communities of Aboriginal women</li> <li>● Create brochures or other educational materials that reflect the lives of Aboriginal women</li> </ul>
Legacy of paternalism	<ul style="list-style-type: none"> <li>● Educate women from a young age about the importance of cervical cancer screening and the importance of the HPV vaccine</li> <li>● Create brochures or other educational materials that reflect the lives of Aboriginal women</li> </ul>
Privacy and confidentiality	<ul style="list-style-type: none"> <li>● Bring services to women who live in remote communities</li> </ul>
Pap test not seen as priority	<ul style="list-style-type: none"> <li>● Educate women from a young age about the importance of cervical cancer screening and the importance of the HPV vaccine</li> <li>● Create brochures or other educational materials that reflect the lives of Aboriginal women</li> </ul>
History of sexual abuse	<ul style="list-style-type: none"> <li>● Use of creative technology for hard-to-reach populations, e.g., self-collection</li> </ul>
Availability of health-care providers	<ul style="list-style-type: none"> <li>● Organize a day or week dedicated to cervical cancer screening</li> <li>● Bring services to the women who live in remote communities</li> </ul>
Fear of focusing on cancer	<ul style="list-style-type: none"> <li>● Educate women from a young age about the importance of cervical cancer screening and the importance of the HPV vaccine</li> </ul>



## Conclusion

**Relationship Building.** “Health promotion in Aboriginal communities is less about transfer of knowledge than about transfer of respect and power,” stated one of the women interviewed. Through the twelve weeks of this project, one of the most important lessons I learned was the importance of building partnerships in Aboriginal communities and with Aboriginal agencies and programs. These partnerships must involve a relationship of mutual respect and power-sharing, and the building of these relationships should precede any other intervention efforts.

**Recommended Interventions.** Following relationship building, this report suggests these activities to promote Aboriginal women’s participation in cervical cancer screening: Educate women from a young age about the importance of cervical cancer screening and the importance of the HPV vaccine in preventing cervical cancer; create brochures and other educational material that reflect the lives of Aboriginal women; organize a day or week dedicated to Pap screening; bring services to the women who live in remote communities; offer drop-in appointments; and use creative technology for hard-to-reach populations.

**The Social Determinants of Health.** These have an enormous impact on the health of Aboriginal women in British Columbia. The legacy of colonialism, the impact of residential schools, and the economic hardships resulting from these issues all play roles in producing poor health outcomes for this population. It is important to acknowledge that many Aboriginal women face the challenges of poverty daily. Outreach projects should be designed in ways that both recognize and attempt to overcome those barriers whenever possible.

**Collaboration with Provincial Health Agencies.** Results from this project indicate that individuals and agencies are very receptive to the idea of working together with provincial health agencies to design and carry out projects of mutual concern and benefit.

**Qualitative research** can play a role in informing and supporting the work of provincial health agencies, as demonstrated by the information and recommendations gathered by these interviews. Future collaborations between qualitative researchers and provincial health agencies should be encouraged.

**Knowledge Transfer.** This report was distributed to the sixteen individuals who participated in the interviews, and to the facilitator of the committee meeting at which the surveys were distributed. Additionally, the report’s finding and recommendations will be presented to the CCSP and a poster session will be held at Simon Fraser University. An abstract will be submitted to the annual BCCA conference and to appropriate publications. Presentations will be held at other venues as they become available. These activities will help to ensure that the information and suggestions contained in the report are returned to the community they are intended to benefit.

## Endnotes

- 1 The term “Aboriginal” is used in this paper to refer to any person who self-identifies as being of Aboriginal heritage, whether First Nations, Métis, or Inuit, on-reserve or off-reserve.
- 2 Thanks to Marina Morrow, Lisa Kan, Anne McCulloch, Dirk Van Niekirk, Greg Hislop, and Annette Browne. I am also deeply grateful to the sixteen women who participated in the interviews.
- 3 Editor’s note: Triangulation refers to the utilization of different data sources to substantiate findings. Where different information sources lead us to the same conclusion, we can be more assured we have made a sustainable finding.
- 4 Editor’s note: While we realize the sample is very small, we believe the insights gained through this study are quite useful.

## References

- British Columbia Cancer Agency. 2007. 2007 Annual Report: Cervical Cancer Screening Program. <[www.bccancer.bc.ca/PPI/Screening/Cervical/about.htm](http://www.bccancer.bc.ca/PPI/Screening/Cervical/about.htm)> Accessed March 10, 2009.
- Browne, A., J. Fiske, and G. Thomas. 2000. “First Nations Women’s Encounters with Mainstream Health Care Services and Systems.” British Columbia Centre of Excellence for Women’s Health.
- Clarke, H., R. Joseph, M. Deschamps, T. Hislop, P. Band, and R. Atleo. 1998. “Reducing cervical cancer among First Nations women.” In *Canadian Nurse* 94(3): 36–41.
- Gandey, A. 2008. “Poverty boosts mortality risk after cancer diagnosis, study shows.” In Medscape Medical News. <[www.medscape.com/viewarticle/576684](http://www.medscape.com/viewarticle/576684)> Accessed June 27, 2008.
- Health Canada. 2004. “Progress Report on Cancer Control in Canada.” <[www.phac-aspc.gc.ca/publicat/prccc/relccc/index.html](http://www.phac-aspc.gc.ca/publicat/prccc/relccc/index.html)> Accessed November 19, 2007.
- Hislop, T., H. Clarke, M. Deschamps, R. Joseph, P. Band, J. Smith, N. Le, and R. Atleo. 1996. “Cervical cytology screening: How can we improve rates among First Nations women in urban British Columbia?” In *Canadian Family Physician* (42): 1701–1708.
- Neuman, W. 2006. *Social research methods: qualitative and quantitative methods*. Boston: Pearson.
- Ogilvie, G., M. Krajden, J. Maginley, J. Isaac-Renton, T. Hislop, R. Elwood-Martin, C. Sherlock, D. Taylor, and M. Rekart. 2007. “Feasibility of self-collection of specimens for human papillomavirus testing in hard-to-reach women.” In *Canadian Medical Association Journal* 177(5): 480–483.
- Pakula, B. 2006. “Access to Cervical Cancer Screening Among First Nations Women and Other Vulnerable Populations in Vancouver’s Downtown Eastside.” Master’s thesis, Simon Fraser University.
- Papps, E., and I. Ramsden. 1996. “Cultural safety in nursing: The New Zealand experience.” In *International Journal for Quality in Health Care* 8(5): 491–497.
- Raphael, D. (ed). 2008. *Social Determinants of Health: Canadian Perspectives*, 2<sup>nd</sup> ed. Toronto: Canadian Scholars’ Press.